Otanes, F. Q. 1922. Head and mouthparts of Mecoptera. Ann. Entomol. Soc. Am. 25:310-323.

Penny, N. D. 1975. Evolution of the extant Mecoptera. J. Kans. Entomol. Soc. 48:331-350.

. 1977. A systematic study of the family Boreidae (Mecoptera). Univ. Kans. Sci. Bull. 51:141–217.

Svensson, S. A. 1972. Boreus Latreille, 1825 (Mecoptera). A synopsis of described species. Entomol. Scand. 3:26–32.

PROC. ENTOMOL. SOC. WASH. 81(1), 1979, p. 31

NOTE

Pinnaspis caricis, New Synonym of Pinnaspis aspidistrae (HOMOPTERA: DIASPIDIDAE)

Pinnaspis caricis Ferris (1957. Proc. Hawaii. Entomol. Soc. 16(2):212) was described from specimens collected on *Ophiopogon japonicus* (Thunb.) Ker-Gawl. (=*Mondo japonica* (Thunb.) Farw.) at Honolulu, Hawaii. According to the description, the median lobes are fused into a single lobe. Because of this character, Ferris compared the species with *Pinnaspis uniloba* (Kuwana) which has fused median lobes but does not otherwise closely resemble *P. caricis*. Ferris also stated that the male of *P. caricis* is not known.

In the type-specimens of *P. caricis* that I examined, the median lobes are not fused but are closely appressed. *Pinnaspis caricis* is identical with *Pinnaspis ophiopogonis* Takahashi (1952. Misc. Rpt. Res. Inst. Nat. Resources Japan. 27:11), which is currently considered a junior synonym of *Pinnaspis aspidistrae* (Signoret) (1869. Ann. Entomol. Soc. Fr. (ser. 4) 9:443) (Takagi. 1970. Insecta Matsumurana. 33(1):106). Although *P. caricis* and *P. ophiopogonis* are apparently parthenogenetic and restricted to few host-plants such as *Liriope* sp., *Ophiopogon* sp. and *Rohdea* sp., there is no reliable morphological character to differentiate them from *P. aspidistrae*. Therefore, until a reliable morphological distinction is found, I think *P. caricis* is best treated as another junior synonym of *P. aspidistrae* (NEW SYN-ONYM).

I thank Dr. S. Takagi, Hokkaido University, Sapporo, Japan, and Mr. R. Schuster, University of California, Davis, for the loan of types and other specimens examined in this study.

Sueo Nakahara, Plant Protection and Quarantine, Animal and Plant Health Inspection Service, USDA, Beltsville, Maryland 20705.